

<p>During Zepto capsulotomy, any OVD with a viscosity of $\leq 300,000$ mPas may be used. Examples:</p>	<p>Viscosity (mPas)</p>
<p>Cohesive Healon Pro Provisc Amvisc Amvisc Plus</p>	<p>150 K 280 K 100 K 60 K</p>
<p>Viscous dispersive DisCoVisc</p>	<p>100 K</p>
<p>Dispersive Healon Endocoat Viscoat NuVisc Cellugel Vitrax</p>	<p>50 K 50 K 45 K 40 K 25 K</p>
<p>Dispersive very low viscosity i-Cell Ocuvis Ocucoat Hymecel Adatocel Visilon</p>	<p>6 K 4.3 K 4 K 4 K 4 K 4 K</p>

<p>After Zepto capsulotomy, any OVD may be used, including those with a viscosity of $> 300,000$ mPas. Examples:</p>	<p>Viscosity (mPas)</p>
<p>Viscoadaptives Healon 5 Pro Healon GV Pro MicroVisc (iVisc) Plus</p>	<p>4 M 2 M 3.3 M</p>
<p>Viscous-cohesive viscoelastic MicroVisc (iVisc) Phaco MicroVisc (iVisc)</p>	<p>1 M 1 M</p>